

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@helloworldbio.com



DATASHEET

ω -Conotoxin GVIA

Product overview

Name	ω -Conotoxin GVIA
Cat No	HB1218
Alternative names	ω -CTX, ω -CgTx; SNX-124
Biological action	Blocker
Purity	>98%
Net peptide content	73% (this peptide is sold by gross weight)
Special requirements	As this product is a toxin, customers are required to complete a short end user declaration when ordering. Our customer care team will be happy to help you with this.
Description	This is a home office notifiable schedule 5 toxin. Potent, selective Ca _v 2.2 (N-type) channel blocker

Biological Data

Biological description	Conotoxin GVIA is a peptide neurotoxin which is a potent, selective and partially reversible Ca _v 2.2 (N-type) calcium channel blocker (IC ₅₀ = 0.15 nM). ω -Conotoxin GVIA binds the Ca _v 2.2 α 1 subunit (α 1B). Conotoxin GVIA inhibits presynaptic Ca _v 2.2 channels, reduces glutamate release and irreversibly blocks synaptic transmission. It also attenuates (RS)-3,5-DHPG-induced LTD (long term depression). It shows antinociceptive action and attenuates hyperalgesia and allodynia. Conotoxin GVIA is active in vivo.
-------------------------------	--

Solubility & Handling

Storage instructions	-20 °C (desiccate)
Solubility overview	Soluble in water (1 mg/ml)
Important	This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

Chemical Data

Chemical name	H-Cys(1)-Lys-Ser-Hyp-Gly-Ser-Ser-Cys(2)-Ser-Hyp-Thr-Ser-Tyr-Asn-Cys(3)-Cys(1)-Arg-Ser-Cys(2)-Asn-Hyp-Tyr-Thr-Lys-Arg-Cys(3)-Tyr-NH ₂ trifluoroacetate salt
Molecular Weight	3037
Molecular Formula	C ₁₂₀ H ₁₈₂ N ₃₆ O ₄₃ S ₆
CAS Number	106375-28-4
PubChem identifier	16133838
SMILES	<chem>C[C@H]([C@H]1C(=O)N[C@H](C(=O)N[C@H](C(=O)N[C@H](C(=O)N[C@H]2C(=O)N[C@H](NC(=O)[C@@H](NC(=O)[C@@H](NC(=O)[C@@H](NC(=O)[C@@H]3C[C@H](CN3C(=O)[C@@H](NC(=O)[C@@H]4C(=O)N[C@H](C(=O)N5C[C@H](C[C@H]5C(=O)N1)O)CO)NC(=O)[C@@H](NC(=O)[C@@H](NC(=O)CNC(=O)[C@@H]6C[C@H](CN6C(=O)[C@@H</chem>

](NC(=O)[C@@H](NC(=O)[C@H](CSSC[C@@H](C(=O)N[C@H](C(=O)N[C@H](C(=O)N4)CO)CCC
NC(=N)N)NC2=O)N)CCCCN)CO)O)CO)CO)CC(=O)N)O)CC7=CC=C(C=C7)O)[C@@H](C)O)CCCC
N)CCCNC(=N)N)C(=O)N[C@@H](CC8=CC=C(C=C8)O)C(=O)N)CC(=O)N)CC9=CC=C(C=C9)O)C
O)O

InChi

InChI=1S/C120H182N38O43S6/c1-53(165)91-114(197)138-66(10-4-6-26-122)95(178)136-68(12-8-
28-132-120(129)130)98(181)149-81(107(190)139-69(93(126)176)29-55-13-19-58(167)20-14-55)49-
204-207-52-84-110(193)153-80-48-203-202-47-64(123)94(177)135-65(9-3-5-25-121)9

InChiKey

FDQZTPPHJRQRQQ-NZPQQUJLSA-N

MDL number

MFCD00076620

References

Complex structures between the N-type calcium channel (CaV2.2) and ω -conotoxin GVIA predicted via molecular dynamics.

Chen R *et al* (2013) *Biochemistry* 52(21)

PubMedID [23651160](#)

Distinct mechanisms contribute to agonist and synaptically induced metabotropic glutamate receptor long-term depression.

Connelly *et al* (2011) *Eur J Pharmacol* 667(1-3)

PubMedID [21575629](#)

The effect of omega-conotoxin GVIA on synaptic transmission within the nucleus accumbens and hippocampus of the rat in vitro.

Home and Kemp (1991) *Br J Pharmacol* 103(3)

PubMedID [1657265](#)

omega-Conotoxin reduces facilitation of transmitter release at the frog neuromuscular junction.

Zengel *et al* (1993) *Brain Res* 611(1)

PubMedID [8100174](#)
