Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET AMN 082 dihydrochloride

Product overview

NameAMN 082 dihydrochlorideCat NoHB0113Biological actionAgonistPurity>98%DescriptionSelective mGlu7 agonist

Images



Biological Data

Biological description	AMN 082 dihydrochloride is a selective mGlu ₇ positive allosteric agonist. It is orally active and blood- brain barrier permeable.
	AMN 082 binds at the mGlu ₇ allosteric site and at \leq 10 μ M fails to show appreciable activating or inhibitory effects at other mGluR subtypes and selected iGluRs.
	Uses
	AMN 082 effects may differ depending on brain region. For example, AMN082 decreases GABA and increases glutamate levels in the nucleus accumbens and amygdala and decreases glutamate and GABA release in the periaqueductal gray (PAG).
	It potently inhibits accumulation of cAMP and stimulates GTP γ S binding (EC ₅₀ = 64 - 290 nM) with agonist efficacies comparable with those of L-AP4. AMN082 also increase splasma corticosterone and ACTH levels.
Application notes	AMN082 reduces fear acquisition and LTP in the amygdala but improves fear extinction. AMN 082 also produces anxiogenic- and anxiolytic-like effects, can facilitate nociception, shows anti-depressant-like activity, reduces ethanol and cocaine intake and facilitates extinction of aversive memories. In vitro use – guidelines
	Non-specific actions may be observed at concentrations of 3-10 μM and above.
	Therefore, for researchers wishing to investigate selective mGlu ₇ actions, it is recommended that this product is not used above concentrations of 1μ M.

<u>In vivo use – guidelines</u>

Guidelines for maximally tolerated doses in vivo are: 6 mg/kg p.o. in mice and 20 mg/kg p.o. in rats

- Those doses result in mGlu₇-dependent physiological effects, e.g. modulation of stresshormones.
- However, non-selective effects have been observed at higher doses (2-3 times higher than those stated above).
- Examples of such non-selective effects include head twists and tremor observed in mGlu₇+/+ (wild-types) and mGlu₇-/- mice (knock-outs).
- The product can be orally administered (p.o.) in a methylcellulose suspension. For further details contact Dr. John F. Cryan at University College Cork. There is currently no data available on maximally tolerated doses for i.v., i.c.v., or i.p. routes of administration.

Use of knock-outs for validation of data

- Dr. Peter J. Flor and his colleagues recommend that the physiological and pharmacological effects of AMN082 should ideally be confirmed by evaluation in mGlu₇+/+ (wild-types) versus mGlu₇-/- mice (KO).
- Effects of AMN082 that are seen in mGlu₇+/+ (wild-types) but not in mGlu₇-/- mice (KO) are most likely mGlu₇+/+ -mediated.
- For details on obtaining mGlu₇+/+ (wild-types) and mGlu₇-/- mice (KO) please contact Dr. Peter J. Flor or Dr. Herman van der Putten.

Solubility & Handling

Storage instructions	+4°C (desiccate)
Solubility overview	Soluble in DMSO (100mM) or water (2mM, gentle warming). Stock solutions (up to 10 mM) can be prepared in DMSO or methanol.
Handling	 If possible, make up solutions and use immediately. If storage of solutions is required, you should aliquot out the solution into tightly sealed vials and store at -20°C and store these for up to one month. Allow the product to equilibrate to RT for at least one hour before opening and using. Always check that your product is completely dissolved before use. The solution should be precipitate-free, clear and colourless. Once removed from -20°C, and brought to room temperature, it is recommended that the solution is used immediately.
Important	This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.

Chemical Data

Chemical name N,N-Bis(diphenylmethyl)-1,2-ethanediamine dihydrochloride Molecular Weight 465.45 **Chemical structure** 2HCI **Molecular Formula** C28H28N2.2HCI **CAS Number** 97075-46-2 **PubChem identifier** 11698390 SMILES C1=CC=C(C=C1)C(C2=CC=C2)NCCNC(C3=CC=CC=C3)C4=CC=CC=C4.CI.CI Synthetic Source InChi InChI=1S/C28H28N2.2CIH/c1-5-13-23(14-6-1)27(24-15-7-2-8-16-24)29-21-22-30-28(25-17-9-3-10-1 8-25)26-19-11-4-12-20-26;;/h1-20,27-30H,21-22H2;2*1H YRQCDCNQANSUPB-UHFFFAOYSA-N InChiKey Appearance White solid

References

Activation of the mGlu7 receptor elicits antidepressant-like effects in mice.

A selective metabotropic glutamate receptor 7 agonist: activation of receptor signaling via an allosteric site modulates stress parameters in vivo.

Mitsukawa K *et al* (2005) Proc Natl Acad Sci U S A 102(51) **PubMedID** 16339898

Metabotropic glutamate 7 receptor subtype modulates motor symptoms in rodent models of Parkinson's disease.

Greco B *et al* (2010) J Pharmacol Exp Ther 332(3) **PubMedID** 19940105