

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

Rapamycin

Product overview

Name	Rapamycin
Cat No	HB2779
Alternative names	Sirolimus
Antibiotic Spectrum	yeast and fungi
Purity	>98%
Description	mTOR inhibitor and potent immunosuppressant. Also used in inducible gene editing methods (e.g. CRISPR/Cas9) and promotes hPSC differentiation.

Biological Data

Biological description

Overview

Rapamycin is an mTOR inhibitor and potent immunosuppressant. It complexes with FKBP12 to bind mTORC1 to inhibit mTOR activity and block subsequent activation of p70 s6 kinase ($IC_{50} = 50 \text{ pM}$).

Active in vivo.

Uses and applications

Rapamycin shows a variety of biological actions. E.g it induces autophagy and apoptosis.

Promotes hPSC differentiation to blood progenitor and mesendoderm cells.

It can also be used in inducible CRISPR/Cas9 systems to enable inducible gene editing.

Used as a chemical dimerizer in Chemically-inducible dimerization (CID).

As mTOR is a central inflammation regulator, rapamycin has recently been investigated as part of cytokine storm / COVID-19 related research.

Solubility & Handling

Storage instructions Solubility overview Important

-20 °C

Soluble in ethanol (20 mM) and DMSO (50 mM)

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.


Chemical Data

Chemical name

(3S,6R,7E,9R,10R,12R,14S,15E,17E,19E,21S,23S,26R,27R,34aS)-9,10,12,13,14,21,22,23,24,25,26,27,32,33,34,34a-Hexadecahydro-9,27-dihydroxy-3-[(1R)-2-[(1S,3R,4R)-4-hydroxy-3-methoxycyclohexyl]-1-methylethyl]-10,21-dimethoxy-6,8,12,14,20,26-hexamethyl-23,27-epoxy-3H-pyrido[2,1-c][1,4]oxaazacyclohentacontine-1,5,11,28,29(4H,6H,31H)-pentone

Molecular Weight

914.18

Chemical structure**Molecular Formula**C₅₁H₇₉NO₁₃**CAS Number**

53123-88-9

PubChem identifier

5284616

SMILESC[C@@H]1CC[C@H]2C[C@@H](/C(=C/C=C/C=C/[C@H](C[C@H](C=O)[C@@H]([C@@H]/C=C/[C@H](C=O)C[C@H](OC(=O)[C@@H]3CCCCN3C(=O)C(=O)[C@@]1(O2)O)[C@H](C)C[C@@H]4CC[C@H]([C@@H](C4)OC)O)C)/C)O)OC)C)/C)OC**InChiKey**

QFJCIRLUMZQUOT-HPLJQQBZSA-N

MDL number

MFCD00867594

Appearance

White to off-white

References

Rapamycin, a specific inhibitor of the mammalian target of rapamycin, suppresses lymphangiogenesis and lymphatic metastasis.

Kobayashi et al (2007) Cancer Sci 98(5)

PubMedID [17425689](#)

Rapamycins: mechanism of action and cellular resistance.

Huang et al (2003) Cancer Biol Ther 2(3)

PubMedID [12878853](#)

Chemical modulators of autophagy as biological probes and potential therapeutics.

Flemin et al (2011) Nat Chem Biol 7(1)

PubMedID [21164513](#)

A split-Cas9 architecture for inducible genome editing and transcription modulation.

Zetsche et al (2015) Nat Biotechnol 33(2)

PubMedID [25643054](#)

Targeting T-cell senescence and cytokine storm with rapamycin to prevent severe progression in COVID-19

Mahe et al (2020) Clin Immunol. 216

PubMedID [32405269](#)
