Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

Pertussis Toxin

Product overview

Name Pertussis Toxin

Cat No HB4729

Alternative names PTX | PT | Islet-activating protein | Holotoxin

Biological action Activator Purity >98%

Description Catalyzes ADP-ribosylation of the G proteins Ga_i , Ga_o and Ga_t

Biological Data

Biological description Toxin produced by Bordetella pertussis . Catalyzes ADP-ribosylation of the α subunits of the

heterotrimeric $G_{i/o}$ proteins; $G\alpha_i$, $G\alpha_o$ and $G\alpha_t$. Blocks receptor coupling and activation by preventing G

protein heterotrimers from interacting with receptors.

This product is not activated. Cells will activate pertussis toxin in an intact system however activation is required in a cell free system. Incubation with high concentrations of dithiothreitol (DTT) can achieve activation, see Kaslow, et al. (1987) for suggested conditions.

Solubility & Handling

Storage instructions

Handling

+4°C for lyophilised or resuspended (do not freeze)

Use solutions within 1 month, long term storage is not recommended.

Before use the suspension should be gently mixed (not vortexed) to make the suspension uniform. Do

not sterile filter.

Pertussis toxin can be inactivated by boiling for 30mins

Important This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not

for human or veterinary use.

Chemical Data

CAS Number70323-44-3SourceB. pertussisAppearanceLyophilised

Formulation Contains 0.05M sodium phosphate and 0.5M sodium chloride at pH 7.2.

References

Tamura et al (1982) Biochemistry 21(22) **PubMedID**6293544

Structure-activity analysis of the activation of pertussis toxin.

Kaslow et al (1987) Biochemistry 26(1) **PubMedID** 3030399

G(i/o) protein-dependent and -independent actions of Pertussis Toxin (PTX).

Mangmool and Kurose (2011) Toxins (Basel) 3(7) **PubMedID** 22069745