Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

4-Hydroxytamoxifen (E) and (Z) isomers (50:50)

Product overview

Name 4-Hydroxytamoxifen (E) and (Z) isomers (50:50)

Cat No HB6095

Alternative names 4-OHT, Afimoxifene, H6278, 4-HT, 4-OH-TAM, OHT, TAM, 4-Hydroxytamoxifen, tamoxifen, z-4oht

Biological action Activator >98%

Description Synthetic estrogen receptor ligand. Widely used in genome engineering (e.g. CreER/ CRISPR-Cas9).

Biological Data

Biological description

Cell permeable, synthetic estrogen receptor ligand. Widely used in genome engineering. (E) and (Z) isomer mix (50:50).

CreER system:

De facto standard compound in drug-inducible manipulation of CreER recombinase. Allows external temporal control of Cre activity *in vivo*.

May also be used for TRAPing / in the TRAP / TRAP2 system (Targeted recombination in active populations).

CRISPR/Cas9 gene editing:

Activates an inactived Cas9 nuclease (rendered inactive by insertion of a 4-OHT dependent-intein) to reduce off-target CRISPR-mediated gene editing (once bound with 4-OHT, conditionally active Cas9s modify target genomic sites with ~25-fold higher specificity than wild-type Cas9).

Also allows tight, repeated on-off control of the nuclease activity of the 'iCas' Cas9 variant which shows high editing efficiency at multiple loci once bound with 4-OHT.

Pure (Z)-4-hydroxytamoxifen also available.

≥70% Z isomer (remainder primarily E-isomer) 4-Hydroxytamoxifen also available.

Solubility & Handling

Solubility overview Storage instructions Handling Soluble in DMSO (100 mM) and in ethanol (50 mM)

-20°C

Storage of solid

 This compound is light sensitive; exposure to light may affect compound performance. You should therefore store the material in the dark and protect from light.

- Do not store the material in solution; make up solutions and use immediately:
- The compound has been shown to isomerise rapidly in solution in most solvents (particularly solvents with a low dielectric constant). You should therefore make up and use solutions immediately.
- The isomerisation process can be precluded by storage of the compound at -25°C in the dark as a THF solution containing ca. 0.025% BHT. (Katzenellenbogen et al (1982) J. Org. Chem. 47 2387.)

Shipping Conditions Important Stable for ambient temperature shipping. Follow storage instructions on receipt.

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.

Chemical Data

Chemical name Molecular Weight Chemical structure 4-(1-[4-(Dimethylaminoethoxy)phenyl]-2-phenyl-1-butenyl)phenol

387.51

HO CH3

Source Synthetic

InChi InChi=1S/C26H29NO2/c1-4-25(20-8-6-5-7-9-20)26(21-10-14-23(28)15-11-21)22-12-16-24(17-13-22

)29-19-18-27(2)3/h5-17,28H,4,18-19H2,1-3H3/b26-25-

InChiKey TXUZVZSFRXZGTL-QPLCGJKRSA-N

Appearance Off-white solid

References

A monohydroxylated metabolite of tamoxifen with potent antioestrogenic activity.

Jordan et al (1977) J Endocrinol 75(2) **PubMedID**591813

A chemical-inducible CRISPR-Cas9 system for rapid control of genome editing.

Liu et al (2016) Nat Chem Biol 12(11)

PubMedID 27618190

Small molecule-triggered Cas9 protein with improved genome-editing specificity.

Small molecule-triggered Cas9 protein with improved genome-editing specificity. (2015) Nat Chem Biol 11(5)

PubMedID 25848930

Simple and efficient production of (Z)-4-hydroxytamoxifen, a potent estrogen receptor modulator.

Yu and Forman (2003) J Org Chem 68(24) **PubMedID**14629178