

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

TFLLR-NH2


Product overview

Name	TFLLR-NH2
Cat No	HB2921
Biological description	TFLLR-NH2 is a selective PAR ₁ (protease-activated receptor-1) activating peptide. It is commonly used to activate platelets and has also been shown to elicit scratching in mice. Active <i>in vivo</i> .
Alternative names	TFLLR, TF-NH2
Biological action	Activator
Purity	>95%
Description	Selective PAR ₁ activating peptide

Solubility & Handling

Storage instructions	-20°C
Solubility overview	Soluble in water (1 mg/ml)
Important	This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

Chemical Data

Chemical name	TFLLR (modifications: C-terminal amide)
Molecular Weight	647.82
Chemical structure	
Molecular Formula	C ₇₆ H ₁₀₄ N ₁₈ O ₁₉ S ₂
Sequence (one letter)	TFLLR
Modifications	C-terminal amide
CAS Number	197794-83-5
PubChem identifier	10146183
SMILES	<chem>C[C@@H]([C@@H](C(=O)N[C@@H](CC1=CC=CC=C1)C(=O)N[C@@H](CC(C)C)C(=O)N[C@@H](CC(C)C)C(=O)N[C@@H](CCCNC(=O)N)C(=O)N)N)O</chem>
InChIKey	ANAMCEKSRDPIPX-GFGQVAFXSA-N
MDL number	MFCD05663482

References

A role for proteinase-activated receptor-1 in inflammatory bowel diseases.

Vergnolle et al (2004) J Clin Invest. 114(10)

PubMedID [15545995](#)

Activation of platelet protease-activated receptor-1 induces epithelial-mesenchymal transition and chemotaxis of colon cancer cell line SW620.

Jia et al (26718652) Oncol Rep 35(2)

Activation of proteinase-activated receptors induces itch-associated response through histamine-dependent and -independent pathways in mice.

Tsuji et al (2008) J Pharmacol Sci. 108(3)

