

DATASHEET

3,3-Diaminobenzidine (DAB) tetrahydrochloride

Product overview

Name	3,3-Diaminobenzidine (DAB) tetrahydrochloride
Cat No	HB0687
Alternative names	DAB, DAB substrate
Biological description	<u>Overview</u>

DAB tetrahydrochloride is a water soluble form of DAB which is a derivative of benzene and is widely used in the staining of nucleic acids and proteins, particularly in IHC and HC procedures.

DAB is oxidized by hydrogen peroxide in the presence of peroxidases such as HRP. Once oxidized, DAB forms a dark brown insoluble precipitate at the reaction site to allow visualisation of the target.

This precipitate can be chelated with osmium tetroxide and the color produced can be enhanced by the addition of metals (e.g. nickel, copper, silver and cobalt).


Uses & applications

In IHC, DAB acts as a substrate for peroxidase enzymes conjugated to a primary or secondary antibody e.g. HRP-conjugated antibodies.

DAB is also used for peroxisome and mitochondrial COX activity staining and used to detect the presence and distribution of hydrogen peroxide in plants cells.

Biological action	DAB also enhances staining of iron.
Purity	Dyes & stains >98%
Description	Chromogen staining agent widely used in IHC

Images



Biological Data

Application notes	DAB Staining using HRP Conjugated Secondary Antibodies
--------------------------	---

Stock solutions:

- 1% DAB (0.1g DAB, 10ml dH₂O, 3-5 drops of 10M HCl), mix well and ensure solution is an even brown color. This can be frozen and subsequently thawed
- 0.3% H₂O₂ (100µl of 30% H₂O₂, 10ml dH₂O), can be kept for short periods of time but better to make fresh

Working solution:

- Mix a 1:1:20 ratio of 1% DAB: 0.3% H₂O₂ : PBS (e.g. 250µl of 1% DAB solution with 250µL of 0.3% H₂O₂ and 5ml PBS which is sufficient for 100 sections)

Protocol:

1. Following secondary antibody incubation and washing add 50µl of DAB working solution to each section and incubate at room temperature for 10 minutes. During this time the color development can be monitored and the incubation time increased or decreased as required.
2. Rinse slides in running tap water for 5 minutes.
3. Counterstain with **hematoxylin** or proceed straight to dehydration and mounting.

Solubility & Handling


Storage instructions
Solubility overview
Important

-20 °C
Soluble in water (2% by mass)
This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

Chemical Data

Chemical name
Molecular Weight
Chemical structure

3,3'-Diaminobenzidine tetrahydrochloride
360.11



Molecular Formula
CAS Number
PubChem identifier
SMILES
InChiKey
MDL number
Appearance

C₁₂H₁₄N₄·4HCl
868272-85-9
16211746
C1=CC(=C(C=C1C2=CC(=C(C=C2)N)N)N)N.O.Cl.Cl.Cl.Cl
DXWSCXIZIHILNP-UHFFFAOYSA-N
MFCD08273058
Off-white to brown solid

References

3,3'-Diaminobenzidine staining interferes with PCR-based DNA analysis.

Doelle et al (2018) Sci Rep. 8(1)

PubMedID [29352159](#)

Detection of Hydrogen Peroxide by DAB Staining in Arabidopsis Leaves.

Daudi et al (2012) Bio Protoc 2(18)

PubMedID [27390754](#)

Chromogen-based immunohistochemical method for elucidation of the coexpression of two antigens using antibodies from the same species.

Nakata et al (2012) J Histochem Cytochem 60(8)

PubMedID [22610462](#)

Comparison of histological techniques to visualize iron in paraffin-embedded brain tissue of patients with Alzheimer's disease.

van Duijin et al (201 3) J Histochem Cytochem 61(11)
PubMedID [23887894](#)

Comparison of histological techniques to visualize iron in paraffin-embedded brain tissue of patients with Alzheimer's disease.

van Duijin et al (201 3) J Histochem Cytochem 61(11)
PubMedID [23887894](#)
