Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

Propidium Iodide Staining Solution (1mg/ml) in water

Product overview

Name Cat No Alternative names Biological description Propidium Iodide Staining Solution (1mg/ml) in water

HB0821

PΙ

Propidium iodide (PI) is a widely used red-fluorescent intercalating agent that binds and labels nucleic acids. This is a 1mg/ml staining solution.

Propidium iodide is membrane impermeant and is therefore frequently used to selectively identify dead cells and is commonly used in flow cytometry to evaluate cell viability.

Propidium iodide (PI) is often used in flow cytometry, fluorescent microscopy and confocal laser scanning microscopy applications.

Frequently used with Oxazole Yellow (YP1) when staining for apoptotic and necrotic cells as apoptotic cells remain impermeant to propidium iodide but permeable to Oxazole Yellow (YP1).


Once bound to the nucleic acids, its fluorescence is enhanced 20- to 30-fold. Wavelength Maxima:


Excitation ~535nm, Emission ~617nm

Biological action Dyes & stains


Description Red-fluorescent cell viability stain. 1mg/ml staining solution in water.

Images









Biological Data

Application notes

#Protocol 1: Propidium Iodide counterstaining of HEK293T cells

- HEK293T cells were cultured on coverslips in 10% FBS in DMEM and fixed with 4% PFA. Immunocytochemistry was performed following our ICC protocol using an anti-vimentin monoclonal antibody at 1μg/ml.
- Prodidium iodide working solution was prepared consisting of 1μg/ml Propidium Iodide and 10μg/ml RNase A and incubated with cells for 30 minutes at room temperature.
- Following washing with PBS, coverslips were mounted and imaged using a confocal microscope using either a 532nm or 514nm laser for excitation

Solubility & Handling

Storage instructions +4°C

Storage of solutions Prepare and use solutions on the same day if possible. Store solutions at -20 °C for up to one month if

storage is required. Equilibrate to RT and ensure the solution is precipitate free before use.

Handling Protect from light.

Shipping Conditions Stable for ambient temperature shipping. Follow storage instructions on receipt.

Important This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not

for human or veterinary use

Chemical Data

Chemical name 2,7-Diamino-9-phenyl-10 (diethylaminopropyl)-phenanthridium iodide methiodide aqueous solution

InChiKey XJMOSONTPMZWPB-UHFFFAOYSA-M

MDL numberMFCD00011921AppearanceRed/pink solution

Excitation 535 nm **Emission** 617 nm

References

The DNA intercalators ethidium bromide and propidium iodide also bind to core histones.

Banerjee A *et al* (2014) FEBS Open Bio 4 **PubMedID** 24649406

Analysis of apoptosis by propidium iodide staining and flow cytometry.

Riccardi C *et al* (2006) Nat Protoc 1(3) **PubMedID**17406435

DNA staining for fluorescence and laser confocal microscopy.

Suzuki T *et al* (1997) J Histochem Cytochem 45(1) **PubMedID** 9010468