

Hello Bio, Inc.  
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500  
F. 609-228-4994

customercare-usa@hellobio.com



## DATASHEET

### Oxazole Yellow Iodide

## Product overview

<b>Name</b>	Oxazole Yellow Iodide
<b>Cat No</b>	HB6210
<b>Alternative names</b>	Yo-Pro-1, YP1, YP-1
<b>Biological description</b>	<b><u>Overview</u></b>

Oxazole Yellow iodide is also known as Yo-Pro-1 or YP1. It is a carbocyanine nucleic acid stain which has a strong binding affinity to nucleic acids.

It is a green fluorescent DNA marker which is commonly used to identify apoptotic cells.

#### **Uses and applications**

Oxazole Yellow (YP1) does not penetrate the plasma membrane of viable cells. However, during apoptosis, apoptotic processes cause the cell-membrane to become slightly permeable. This allows Oxazole Yellow (YP1) to enter these cells and bind to nucleic acids to allow detection of apoptotic cells.

It is frequently used with **propidium iodide** when staining for apoptotic and necrotic cells as apoptotic cells remain impermeant to propidium iodide but permeable to Oxazole Yellow (YP1).

<b>Biological action</b>
<b>Purity</b>
<b>Description</b>

Dyes & stains

>99%

Oxazole Yellow iodide is also known as Yo-Pro-1. It is a commonly used apoptosis marker.

## Biological Data


<b>Application notes</b>	Oxazole yellow (also known as YO-PRO1) is a cyanine dye with strong affinity for nucleic acids used to stain apoptotic cells. Oxazole yellow from Hello Bio has an excitation peak at 491nm (when recorded at 507nm) and peak emission at 505nm (when excited at 490nm). For protocol see #Protocol 1 in application notes below.
--------------------------	---

## Solubility & Handling

<b>Storage instructions</b>	-20°C (protect from light)
<b>Solubility overview</b>	Soluble in water (1 mg/ml), and in DMSO (1 mg/ml)
<b>Important</b>	This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

## Chemical Data

<b>Chemical name</b>	4-[(3-Methyl-2(3H)-benzoxazolylidene)methyl]-1-[3-(trimethylammonio)propyl]-quinolinium diiodide
<b>Molecular Weight</b>	629.32
<b>Chemical structure</b>	



<b>Molecular Formula</b>	C <sub>24</sub> H <sub>29</sub> I <sub>2</sub> N <sub>3</sub> O
<b>CAS Number</b>	152068-09-2
<b>PubChem identifier</b>	6439500
<b>SMILES</b>	CN\1C2=CC=CC=C2O/C1=C\C3=CC=[N+](C4=CC=CC=C34)CCC[N+](C)(C)C.[I-].[I-]
<b>Source</b>	Synthetic
<b>InChiKey</b>	ULHRKLSNHXXJLO-UHFFFAOYSA-L
<b>Appearance</b>	Yellow solid

---

## References

### Application of the novel nucleic acid dyes YOYO-1, YO-PRO-1, and PicoGreen for flow cytometric analysis of marine prokaryotes

Marie D *et al* (1996) Appl Environ Microbiol 62(5)

**PubMedID** [8633863](#)

### Rapid quantification of cell viability and apoptosis in B-cell lymphoma cultures using cyanine SYTO probes

Wlodkowic D *et al* (2011) Methods Mol Biol 740

**PubMedID** [21468970](#)

### Evaluation of YO-PRO-1 as an early marker of apoptosis following radiofrequency ablation of colon cancer liver metastases

Fujisawa S *et al* (2014) Cytotechnology 66(2)

**PubMedID** [24065619](#)

### Evaluation of a continuous quantification method of apoptosis and necrosis in tissue cultures

Gawlitta D *et al* (2004) Cytotechnology 46(2-3)

**PubMedID** [19003268](#)

---