



DATASHEET

Protease Inhibitor Cocktail solution (mammalian cell and tissue extract use) (EDTA free, 100x in DMSO)

Product overview

Name	Protease Inhibitor Cocktail solution (mammalian cell and tissue extract use) (EDTA free, 100x in DMSO)
Cat No	HB9081
Description	Protease Inhibitor Cocktail (EDTA free) solution for mammalian cell and tissue extract use

Biological Data

Biological description

Overview

This protease inhibitor cocktail contains a mixture of protease inhibitors with broad specificity for various proteases. It is optimized for use in mammalian cell and tissue extracts. Protease inhibitor cocktails protect proteins from degradation by endogenous proteases released during protein extraction and purification. This cocktail is EDTA free.

Components and action

The following components are included in this cocktail which target the major classes of proteases including serine, cysteine and acid proteases and also aminopeptidases.

- AEBSF hydrochloride serine protease inhibitor (100mM)
- Aprotinin (bovine recombinant, Nicotiana sp.): serine protease inhibitor (80μM)
- Bestatin: aminopeptidase B and leucine aminopeptidase inhibitor (5mM)
- E-64: cysteine protease inhibitor (1.5mM)
- Leupeptin hemisulfate: serine/cysteine protease inhibitor (2mM)
- Pepstatin A: aspartic protease inhibitor (1mM)

Formulation & usage recommendation

The cocktail is supplied as a 1ml solution of 100X concentrate in DMSO. It is generally effective at a 1X final concentration but may require optimization if a sample contains particularly high levels of proteases.

1mL of this cocktail solution is recommended for inhibition of endogenous enzymes found in 100 mL of lysate from 20g of bovine liver or 10 mL of lysate from CHO cells. Please note that not all lysates contain the same levels of endogenous enzymes and you may need to adjust the volume of cocktail required accordingly.

Application notes

1. Equilibrate the bottle to room temperature.
2. Vortex the bottle to ensure a homogeneous suspension.
3. Immediately before use, add 10μL of the Protease Inhibitor Cocktail per milliliter of sample directly to the lysis buffer or extract to produce a 1X final concentration.

Solubility & Handling

Storage instructions	-20 °C
Solubility overview	DMSO (supplied in 1ml DMSO)

Handling	This cocktail may freeze during storage. This does not affect product performance or function.
Important	Equilibrate the bottle to room temperature until the cocktail becomes a liquid before use. This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.

Chemical Data

Appearance	Brown liquid
------------	--------------
