





DATASHEET

Janelia Fluor® 549, SE

Product overview

Name	Janelia Fluor® 549, SE
Cat No	HB7336
Biological description	Cell permeable, yellow fluorescent dye supplied as an NHS ester / succinimidyl ester (SE) for coupling to primary amine (NHS) groups. NHS esters are used to label the primary amines of proteins and are commonly used for conjugating dyes to a protein or antibody.
	May be used for cellular imaging when combined with the HaloTag or SNAP-tag self labelling systems. Suitable for confocal microscopy and super resolution microscopy (SRM) including techniques such as dSTORM (both live and fixed cells) and STED. Also suitable for flow cytometry. Janelia Fluor® 549 is 2 x brighter than TMR and Cy3 <i>in vitro</i> and live-cell experiments.
Alternative names	Spectrally similar dyes: Alexa Fluor® 546, Alexa Fluor® 555, BDY TMR-X, Atto 550, CF 555, TAMRA, Cyanine 3
Biological action	JF549, JF549-SE
Description	Dyes & stains Yellow dye for coupling to NHS groups. Suitable for dSTORM, STED, confocal microscopy, live cell imaging and flow cytometry.

Images



Biological Data

Application notes

#Protocol 1: Conjugation of Janelia® Fluor 549 SE to antibodies

- Using a desalting column, perform buffer exchange (following manufacturer instructions) of antibody into a carbonate buffer (100mM, pH 8-8.25)
- Mix together the antibody and Janelia® Fluor 549 SE (prepared at 10mM in anhydrous DMSO or DMF) in a 15:1 molar ratio. Incubate in the dark for 60 minutes at room temperature with gentle mixing.
- Add 10% by volume of 0.75M Tris-HCl pH7.4 (to a final concentration of 75mM) to stop the conjugation reaction. Incubate for 10-15 minutes at room temperature in the dark with gentle mixing.
- Use a desalting column, perform buffer exchange (following manufacture instructions) of antibody into PBS 0.05% sodium azide. This step also removes any unbound dye

#Protocol 2: Immunocytochemistry

- ICC was performed upon 4% PFA fixed HeLa cells using **FITC Phalloidin** (1:500) and a anti- β tubulin monoclonal antibody (HB6491, 1:2,000 / 0.5 μ g/ml). A polyclonal goat anti-mouse Janelia[®] Fluor 549 conjugated antibody was used at a dilution of 1:300 as a secondary antibody.
- Please see our detailed [immunocytochemistry protocol](#) for details of the full protocol.

#Protocol 3: Immunohistochemistry

- 40 μ m horizontal sections were cut from a 4% PFA fixed rat brain.
- IHC(IF) was performed using mouse monoclonal anti-GFAP (**HB8267**, 1:1000 dilution / 1 μ g/ml) and rabbit monoclonal anti-NFL (**HB7266**, 1:2000 / 0.5 μ g/ml) antibodies. A polyclonal goat anti-mouse Janelia[®] Fluor 549 conjugated antibody was used at a dilution of 1:300 as a secondary antibody.
- Please see our detailed [immunohistochemistry protocol](#) for details of the full protocol

Solubility & Handling

Storage instructions Solubility overview Handling

-20 °C

Soluble in DMSO

This compound is light and temperature sensitive; exposure to light may affect compound performance. We therefore recommend storing the material in a freezer and protecting from light.

Shipping conditions Important


Ship on ice

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

Chemical Data

Chemical name Molecular Weight Chemical structure

3,6-Di-1-azetidiny-9-[2-carboxy-5-[(2,5-dioxo-1-pyrrolidinyloxy)carbonyl]phenyl]xanthylum, inner salt 551.5



Molecular Formula CAS Number PubChem identifier SMILES

C₃₁H₂₅N₃O₇

1811539-32-8

124201856

C1CN(C1)C2=CC3=C(C=C2)C(=C4C=CC(=[N+](C4)C(=O)O)C6=C(C=CC(=C6)C(=O)ON7C(=O)CCC7=O)C(=O)[O-])

Source InChiKey Licensing details

Synthetic

DWUIXAGWHCMXHN-UHFFFAOYSA-N

Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus

References

A general method to improve fluorophores for live-cell and single-molecule microscopy.

Grimm JB et al (2015) Nature methods 12

PubMedID

[25599551](#)

A general method to fine-tune fluorophores for live-cell and in vivo imaging.

Grimm JB et al (2017) Nature methods 14

PubMedID [28869757](#)

Synthesis of Janelia Fluor HaloTag and SNAP-Tag Ligands and Their Use in Cellular Imaging Experiments.

Grimm JB et al (2017) Methods in molecular biology (Clifton, N.J.) 1663

PubMedID [28924668](#)

Rational Design of Bioavailable Photosensitizers for Manipulation and Imaging of Biological Systems.

Binns TC et al (2020) Cell chemical biology 27

PubMedID [32698018](#)
