


## DATASHEET

### Anti-Myelin Basic Protein (MBP) Antibody ValidAb™

#### Product overview

<b>Name</b>	Anti-Myelin Basic Protein (MBP) Antibody ValidAb™
<b>Cat No</b>	HB8014
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Target</b>	Myelin basic protein
<b>Description</b>	Antibody to myelin basic protein (MBP) - marker for oligodendrocytes and Schwann cells. Part of the ValidAb™ range of highly validated, data-rich antibodies.

#### Validation data



#### Product information

<b>Immunogen</b>	Myelin basic protein (MBP) purified from bovine brain
<b>Epitope</b>	Amino acids 145 - 184 of the human 21.5kDa sequence corresponding to the amino acid sequence: AEGQRPGFGYGGRASDYKSAHKGFKGVDAQGTLSKIFKLG
<b>Clone number</b>	7D2
<b>Isotype</b>	IgG1
<b>Purification</b>	Protein G affinity purification
<b>Concentration</b>	1 mg/ml
<b>Formulation</b>	50% PBS, 50% glycerol + 5mM sodium azide
<b>Predicted species reactivity</b>	Rat, Human, Pig, Horse, Cow
<b>Tested species reactivity</b>	Rat, Mouse (no staining)

#### Tested applications

<b>Applications</b>	WB, IHC(IF)
<b>Western blot optimal concentration</b>	250ng/ml (1:4,000 dilution) as tested in a rat brain P2 membrane preparation
<b>IHC(IF) optimal concentration</b>	500ng/ml (1:2,000 dilution) as tested in rat cerebellum sections
<b>Positive control</b>	Myelin basic protein is present in large quantities within the CNS and PNS therefore brain and/or other nerve samples form an excellent positive control. MBP expression has been reported (see the <a href="#">human protein atlas</a> ) in some cell lines such as SK-MEL-30 cells.
<b>Negative control</b>	MBP is not found in appreciable quantities in peripheral tissues therefore these can be used as a negative control. Within the brain MBP is associated with the membrane bound fraction and is absent from the cytosol therefore this can be used as a negative control. MBP is also absent from many common cells lines such as SH-SY5Y, HeLa and HEK293 cells.
<b>Open data link</b>	Please follow this <a href="#">link to OSF</a>

## Target information

<b>Other names</b>	Myelin A1 protein, Myelin membrane encephalitogenic protein
<b>UniProt ID</b>	P02686
<b>Gene name</b>	MBP
<b>NCBI full gene name</b>	myelin basic protein
<b>Entrez gene ID</b>	4155
<b>Amino acids</b>	160 - 304 (17.3 - 33.1kDa) depending upon isotype
<b>Isoforms</b>	Myelin basic protein has a number of isoforms expressed under the control of alternative splicing: <ul style="list-style-type: none"> <li>• Isoform 1 (canonical), also known as Golli-MBP1, HOG7: 304aa, 33.1kDa</li> <li>• Isoform 2, also known as Golli-MBP2, HOG5: 197aa, 21.5kDa</li> <li>• Isoform 3, also known as MBP1, 197aa, 21.5kDa</li> <li>• Isoform 4, also known as MBP2, 186aa, 20.2kDa</li> <li>• Isoform 5, also known as MBP3, 171aa, 18.6kDa</li> <li>• Isoform 6, also known as MBP4, 160 aa, 17.3kDa</li> </ul>
<b>Expression</b>	MBP isoforms are expressed widely in the CNS and PNS within the myelin sheaths that surround axons. Oligodendrocytes in the CNS and their equivalent in the PNS, Schwann cells, express MBP strongly therefore MBP is a good marker for them. The golli forms of MBP are also expressed in the immune system and bone marrow.
<b>Subcellular expression</b>	Expressed within the cytosol of oligodendrocytes and Schwann cells and the myelin sheath of axons. The 21.kDa isoform (MBP1) is also found in the nucleus of oligodendrocytes.
<b>Processing</b>	The initiator methionine is removed from isoforms 3-6.
<b>Post translational modifications</b>	MBP isoforms are subject to numerous post-translational modifications including phosphorylation, citrullination and acetylation. Some of these modification fall within the epitope of HB8014.
<b>Homology (compared to human)</b>	Mouse and rat isoforms have a 74.6% and 92.9% identity to human MBP in a BLAST search
<b>Similar proteins</b>	No similar proteins reported in a BLAST search
<b>Epitope homology (between species)</b>	In a BLAST search only MBP resulted as a match with the epitope sequence.
<b>Epitope homology (other proteins)</b>	In a BLAST search the following species' MBP proteins had the following homology with the epitope sequence: <ul style="list-style-type: none"> <li>• Human – 100% identity</li> <li>• Bovine – 92.5% identity</li> <li>• Chimpanzee – 95.1% identity</li> <li>• Rat – 92.7% identity</li> <li>• Mouse – 92.7% mouse</li> <li>• Rabbit – 85% identity</li> <li>• Horse – 85.4% identity</li> <li>• Pig – 85.4% identity</li> <li>• Chicken – 61.0% identity</li> </ul>

## Storage & Handling

<b>Storage instructions</b>	-20°C
<b>Important</b>	This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

## References

### **Myelin basic protein: a multifunctional protein.**

Boggs JM (2006) Cellular and molecular life sciences : CMLS 63

**PubMedID** [16794783](#)

### **Multiple sclerosis and myelin basic protein: insights into protein disorder and disease.**

Martinsen V et al (2022) Amino acids 54

**PubMedID** [34889995](#)

### **Myelin basic protein immunoreactivity in the human embryonic CNS**

Zecevic N et al (1998) Brain research. Developmental brain research 105

**PubMedID** [9473608](#)

### **Immunosignals of Oligodendrocyte Markers and Myelin-Associated Proteins Are Critically Affected after Experimental Stroke in Wild-Type and Alzheimer Modeling Mice of Different Ages.**

Michalski D et al (2018) Frontiers in cellular neuroscience 12

**PubMedID** [29467621](#)

### **The myelin basic protein gene is expressed in differentiated blood cell lineages and in hemopoietic progenitors.**

Marty MC et al (2002) Proceedings of the National Academy of Sciences of the United States of America 99

**PubMedID** [12084930](#)

### **gamma-Aminobutyric acid outside the mammalian brain.**

Erdö SL et al (1990) Journal of neurochemistry 54

**PubMedID** [2405103](#)

---