

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@hellobio.com




DATASHEET

Anti-BrdU antibody ValidAb™

Product overview

Name	Anti-BrdU antibody ValidAb™
Cat No	HB9919
Host	Mouse
Clonality	Monoclonal
Target	BrdU
Description	Antibody to BrdU - thymidine analogue incorporated into DNA during replication therefore used as a marker of proliferating cells. Part of the ValidAb™ range of highly validated, data-rich antibodies.

Validation data



Product information

Immunogen	BrdU conjugated with hemocyanine.
Clone number	MoBu-1
Isotype	IgG1
Purification	Protein A affinity chromatography
Formulation	Lyophilised. When reconstituted contains PBS with 15mM sodium azide and 1% recombinant albumin
Predicted species reactivity	NA
Tested species reactivity	NA

Tested applications

Applications	ICC, IHC(IF)
IHC(IF) optimal concentration	1 µg/ml (1:1000) as measured in rat hippocampus.

ICC optimal concentration
Product specific protocols

1 µg/ml (1:1000) as measured in mixed neuronal cell cultures.
The dense structure of chromatin can prevent anti-BrdU antibodies binding to the intercalated BrdU within the DNA helix. Denaturing the DNA can therefore improve staining:

- Incubate brain sections or coverslips in 2M HCl for 30 minutes at 37 °C
- Incubate with 0.1M sodium tetraborate (2 x 5 minute incubations) to neutralise the acid
- Wash in PBS / TBS (3 x 5 minute washes)
- Continue with immunostaining (see our [IHC\(IF\)](#) and [ICC](#) protocols for more information)

Positive control
Negative control
Open data link

For more details on BrdU immunostaining please see [Wojtowicz and Kee., 2006](#)
Any cell line or tissue that has had BrdU administered to it while cells are replicating
Any cell line or tissue that has not been exposed to BrdU
Please follow [this link](#) to the OSF.

Target information

Other names [5-Bromo-2-deoxyuridine](#)

Storage & Handling

Storage instructions -20 °C then use reconstitution advice
Reconstitution advice We recommend reconstituting with either:

- dH₂O and storing at 4 °C
- 50:50 ratio of dH₂O to glycerol and storing at -20 °C
- dH₂O then aliquot and store at -80 °C

Take care when opening as the precipitate is extremely light and can easily be lost if disturbed. When reconstituting make sure that the antibody is thoroughly dissolved by pipetting up and down before giving the antibody a brief spin at <10,000g to make sure that all material is recovered and at the bottom of the tube.

Important

For more information please see our detailed guide on [storing and using your antibody](#)
This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

References

[BrdU assay for neurogenesis in rodents.](#)

Wojtowicz JM et al (2006) Nature protocols 1
PubMedID [17406427](#)

[The use of bromodeoxyuridine incorporation assays to assess corneal stem cell proliferation.](#)

Crane AM et al (2013) Methods in molecular biology (Clifton, N.J.) 1014
PubMedID [23690005](#)

[Proliferation assays \(BrdU and EdU\) on skeletal tissue sections.](#)

Mead TJ et al (2014) Methods in molecular biology (Clifton, N.J.) 1130
PubMedID [24482177](#)

[Neurogenesis in the adult human hippocampus.](#)

Eriksson PS et al (1998) Nature medicine 4
PubMedID [9809557](#)
