Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



DATASHEET

Anti-NeuN antibody ValidAb™

Product overview


Name Anti-NeuN antibody ValidAbTM


Cat NoHB6498Alternative namesFox-3HostRabbitClonalityPolyclonalTargetNeuN


Description Antibody to NeuN - marker for mature neurones expressed in the nucleus. Part of the ValidAb™ range


of highly validated, data-rich antibodies.


Validation data























Product information

Epitope Amino acids 5-24 of human NeuN

Isotype IgG

Purification Immunogen affinity purification

Concentration 1mg/ml

Formulation 50% PBS, 50% glycerol + 5mM sodium azide

Predicted species reactivity Mouse, Rat, Human Tested species reactivity Mouse, Rat

Tested applications

Applications WB, IHC(IF)

Western blot optimal 0.5μg/ml (1:2000 dilution) as tested in a rat brain cytosol preparation

concentration

IHC(IF) optimal concentration 0.5μg/ml (1:2000 dilution) as tested in rat brain sections

Positive control NeuN is highly expressed in the neurons of the CNS and PNS. It is also expressed in SH-SY5Y cells.

Negative controlAny tissue not of neural origin. Most cell lines are NeuN negative.

Open data link Please follow this link to OSF

Target information

Other names FOX3, RNA binding protein fox-1 homolog 3, Fox-1 homolog C, RBFOX3, RFOX3

UniProt ID A6NFN3
Gene name RBFOX3

NCBI full gene name RNA binding fox-1 homolog 3

Entrez gene ID 146713

Amino acids Dependent on isoform

Isoforms NeuN binds primarily to FOX3 which has two isoforms. Isoform 1 is described as the canonical

sequence with 312 amino acids (33.8kDa) while isoform 2 has a 13 residue insert at position 312 leading to a total length of 325 amino acids (35.1kDa). NeuN antibodies also bind to synapsin-1 in western blot experiments (but not in IHC or ICC) which has two isoforms. Isoform 1 is 705aa long

(74.1kDa) while isoform 2 is shorter at 669aa (70.0kDa).

Expression NeuN is expressed only within neurones. While the vast majority of neurones express NeuN some cell

types such as Purkinje cells, stellate and golgi cells do not show immunoreactivity.

Subcellular expression Expression is primarily localised to the nucleus however some FOX3 isoforms can localise to the

cytosol.

Processing None

Post translational Phosphorylation has been reported (see Lind et al., 2004. J Neurosci Res. 79: 295-302) which is

modifications directly related to immunoreactivity whereby dephosphorylation abolished staining.

Homology (compared to

human)

Mouse FOX3 shows 95.02% identity to human FOX3 wheras rat FOX3 shows no similarity due to a

large 47 residue insertion at amino acid 252 in rats.

Similar proteins RNA-binding protein fox-1 homolog 1 (40-44kDa) shows 67.3% identity while RNA-binding protein

fox-1 homolog 2 (37-47kDa) shows 56.5% identity

Storage & Handling

Storage instructions

-20°C

Important

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not

for human or veterinary use

References

NeuN: a useful neuronal marker for diagnostic histopathology.

Wolf HK et al (1996) The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society 44

PubMedID 8813082

NeuN As a Neuronal Nuclear Antigen and Neuron Differentiation Marker.

Gusel'nikova VV et al (2015) Acta naturae 7 **PubMedID** 26085943

Identification of neuronal nuclei (NeuN) as Fox-3, a new member of the Fox-1 gene family of splicing factors.

Kim KK et al (2009) The Journal of biological chemistry 284

PubMedID 19713214

Characterization of the neuronal marker NeuN as a multiply phosphorylated antigen with discrete subcellular localization.

Lind D et al (2005) Journal of neuroscience research 79

PubMedID 15605376

Novel Insights into NeuN: from Neuronal Marker to Splicing Regulator.

Duan W et al (2016) Molecular neurobiology 53 **PubMedID** 25680637