


DATASHEET

Anti-GFAP antibody ValidAb™

Product overview

Name	Anti-GFAP antibody ValidAb™
Cat No	HB6406
Host	Chicken
Clonality	Polyclonal
Target	GFAP
Description	Antibody to GFAP - cytoskeletal protein used as an astrocyte marker. Part of the ValidAb™ range of highly validated, data-rich antibodies.

Validation data



Product information

Immunogen	Recombinant human GFAP (isoform 1) expressed in and purified from <i>E. coli</i>
Isotype	IgY
Purification	Unpurified

Formulation	Lyophilised. When reconstituted contains IgY preparation with 5mM sodium azide and 1% recombinant BSA.
Predicted species reactivity	Mouse, Rat, Human, Pig, Horse, Cow
Tested species reactivity	Mouse, Rat

Tested applications

Applications	ICC, WB, IHC(IF)
Western blot optimal concentration	1:8,000 dilution as tested in a rat brain cytosol preparation.
IHC(IF) optimal concentration	1:4,000 dilution as tested in free-floating paraformaldehyde fixed horizontal rat brain sections.
ICC optimal concentration	1:8,000 dilution as tested in cultured rat neurones.
Positive control	GFAP is highly expressed in neural tissues containing astrocytes. It is not widely expressed in cell lines, however it is in specific lines such as U-87 MG.
Negative control	Most non-neural tissues. Please note that GFAP expression has been reported in a subset of pancreatic and hepatic cells in rats and mice kidney cells. It is generally poorly expressed in common cell lines such as HeLa or HEK293.
Open data link	Please follow this link to OSF

Target information

UniProt ID	P14136
Gene name	GFAP
NCBI full gene name	glial fibrillary acidic protein
Entrez gene ID	2670
Amino acids	432 (49.9kDa)
Isoforms	GFAP has three confirmed and 21 potential isoforms. Isoform 1 (GFAP alpha): canonical, 49.9kDa; Isoform 2 (GFAP epsilon): amino acid changes between positions 391 and 432, 49.5kDa; Isoform 3 (GFAP kappa): amino acid changes between positions 391 and 432, 50.3kDa
Expression	GFAP is primarily expressed within astrocytes of the central nervous system alongside also expressing in non-myelinating Schwann cells of the peripheral nervous system and satellite cells of the peripheral ganglia. GFAP expression has also been reported in Leydig cells of the testis alongside stellate cells from the pancreas and liver in rats.
Subcellular expression	GFAP is a key cytoskeletal component therefore is widely expressed as bundles of GFAP positive fibres.
Target function	GFAP is a class III intermediate filament protein with an important role in many processes beyond being a structural cytoskeletal component. GFAP is involved in mitosis of astrocytes, mediating interactions between astrocytes and neurones and repair after CNS injury.
Processing	Following translation, no processing is required for GFAP to reach its active conformation.
Post translational modifications	GFAP is subjected to numerous post-translational modifications including 9 phosphorylation sites which are the target of AURKB and ROCK1 alongside 5 separate citrullination sites.
Homology (compared to human)	Rat, mouse and human GFAP proteins have a 90% similarity score in a direct BLAST comparison.
Similar proteins	Other type III intermediate filament proteins have homology with GFAP including Vimentin (58%), Desmin (59%) and Peripherin (56%) when assessed using BLAST.

Storage & Handling

Storage instructions	-20 °C then use reconstitution advice
Reconstitution advice	Upon receipt store at either -20 °C or -80 °C.

For 100µg packs either:

- Reconstitute with 100µl dH₂O and store at 4 °C
- Reconstitute with 50µl dH₂O and 50µl glycerol then store at -20 °C
- Reconstitute with 100µl dH₂O, aliquot then snap freeze and store at -80 °C

For 25µg packs either:

- Reconstitute with 25µl dH₂O and store at 4 °C
- Reconstitute with 12.5µl dH₂O and 12.5µl glycerol then store at -20 °C
- Reconstitute with 25µl dH₂O, aliquot then snap freeze and store at -80 °C

Important

For more information [read our guide](#) on the best care for your product. Take care when opening as the precipitate is extremely light and can easily be lost if disturbed. When reconstituting make sure that the antibody is thoroughly dissolved by pipetting up and down before giving the antibody a brief spin at 10,000g to make sure that all material is recovered and at the bottom of the tube.

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

References

Glial fibrillary acidic protein: GFAP-thirty-one years (1969-2000).

Eng LF et al (2000) Neurochemical research 25

PubMedID [11059815](#)

GFAP-expressing progenitors are the principal source of constitutive neurogenesis in adult mouse forebrain.

Garcia AD et al (2004) Nature neuroscience 7

PubMedID [15494728](#)

GFAP and astrogliosis.

Eng LF et al (1994) Brain pathology (Zurich, Switzerland) 4

PubMedID [7952264](#)

Importance of GFAP isoform-specific analyses in astrocytoma.

van Bodegraven EJ et al (2019) Glia 67

PubMedID [30667110](#)

The role of GFAP and vimentin in learning and memory.

Wilhelmsson U et al (2019) Biological chemistry 400

PubMedID [31063456](#)
