Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



# DATASHEET

DCFDA / H2DCFDA - Cellular ROS Assay Kit

#### **Product overview**

Name Cat No Biological description DCFDA / H2DCFDA - Cellular ROS Assay Kit HB7375 DCFDA / H<sub>2</sub>DCFDA is a cell permeable fluorescent probe that is redox sensitive and used to measure the concentration of reactive oxygen species (ROS) within a population of cells. DCFDA / H<sub>2</sub>DCFDA diffuses into cells where it is hydrolysed by intracellular esterases into a non-fluorescent and non-cell permeable intermediate. Upon reaction with ROS this forms the fluorescent compound 2',7' -dichlorofluorescein (DCF) which is excited at 485nm and emits at 535nm. Pyocyanin is included within this kit as a positive control. Pyocyanin promotes the formation of ROS through inactivation of catalase and depleting reduced glutathione.

This kit contains:


• DCFDA / H<sub>2</sub>DCFDA assay reagent

Kit for measurement of reactive oxygen species (ROS) within cells.

- DMSO
- Lyophilised Pyocyanin
- 10x assay buffer Cell Culture, FACS and flow cytometry, ICC

Applications Description

## Images



### **Biological Data**

**Application notes** 

Please follow this link to a full DCFDA / H2DCFDA - Cellular ROS Assay Kit protocol

## **Solubility & Handling**

 Storage instructions
 -20°C

 Important
 This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

### **Chemical Data**

**Chemical name** 

Molecular Weight Chemical structure

Molecular Formula CAS Number PubChem identifier SMILES InChiKey 485.27

4091-99-0 77718 CC(=O)OC1=C(C=C2C(C3=CC(=C(C=C3OC2=C1)OC(=O)C)CI)C4=CC=C4C(=O)O)CI PXEZTIWVRVSYOK-UHFFFAOYSA-N

#### References

Detection of Total Reactive Oxygen Species in Adherent Cells by 2',7'-Dichlorodihydrofluorescein Diacetate Staining.

Kim H et al (2020) Journal of visualized experiments : JoVE**PubMedID**32658187

The involvement of TLR2 in cytokine and reactive oxygen species (ROS) production by PBMCs in response to Leishmania major phosphoglycans (PGs).

Kavoosi G et al (2009) Parasitology 136 **PubMedID** 19631014