Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@hellobio.com



## **DATASHEET**

Goat Anti-Mouse IgG H&L (Biotin) preadsorbed ValidAb<sup>TM</sup>

### **Product overview**

Name Goat Anti-Mouse IgG H&L (Biotin) preadsorbed ValidAb<sup>TM</sup>


Cat NoHB11345HostGoatClonalityPolyclonalTargetMouse IgG H&L

**Conjugate** Biotin

**Description** Goat Anti-Mouse IgG H&L (Biotin) preadsorbed secondary antibody. Part of the ValidAb<sup>TM</sup> range of

highly validated, data-rich antibodies.

#### Validation data



## **Product information**

**Immunogen** Purified mouse IgG

Isotype IgG

**Purification** Immunogen affinity chromatography. Pre-adsorbed with human serum proteins

Concentration 0.5 mg/m

Formulation Lyophilised. When reconstituted contains PBS with 1% recombinant albumin and less than 0.1%

sodium azide

## **Tested applications**

**Applications** ELISA, ICC, IHC, IHC-P, WB, IHC(IF)

Western blot optimal Tested at 1:10,000. Optimise dependent upon assay.

concentration

 $\begin{tabular}{ll} \textbf{IHC(IF) optimal concentration} & Tested at 1:250 (2 $\mu g/ml)$. Optimise dependent upon assay. \\ \textbf{IHC-P optimal concentration} & Tested at 1:300 dilution. Optimise dependent upon assay. \\ \textbf{ICC optimal concentration} & Tested at 1:250 (2 $\mu g/ml)$. Optimise dependent upon assay. \\ \end{tabular}$ 

Negative control

While this antibody has been cross-adsorbed to reduce non-specific binding it is still often worthwhile to conduct a control experiment where the primary antibody is omitted to give confidence that the

## **Storage & Handling**

# Storage instructions Reconstitution advice

-20 °C then use reconstitution advice We recommend reconstituting with either:

- dH<sub>2</sub>O and storing at 4°C
- 50:50 ratio of  $dH_2O$  to glycerol and storing at -20 °C
- dH<sub>2</sub>O then aliquot and store at -80°C

Take care when opening as the precipitate is extremely light and can easily be lost if disturbed. When reconstituting make sure that the antibody is thoroughly dissolved by pipetting up and down before giving the antibody a brief spin at <10,000g to make sure that all material is recovered and at the bottom of the tube.

**Shipping Conditions Important** 

For more information please see our detailed guide on storing and using your antibody Stable for ambient temperature shipping. Follow storage instructions on receipt.

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

## References

The use of antiavidin antibody and avidin-biotin-peroxidase complex in immunoperoxidase technics.

Hsu SM et al (1981) American journal of clinical pathology 75

PubMedID 6167159

### **Antigen Retrieval and Signal Amplification.**

Dunkenberger L et al (2022) Methods in molecular biology (Clifton, N.J.) 2422

**PubMedID** 34859399