

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@hellobio.com




DATASHEET

Goat Anti-Rabbit IgG H&L (Biotin) preadsorbed ValidAb™

Product overview

| | |
|--------------------|---|
| Name | Goat Anti-Rabbit IgG H&L (Biotin) preadsorbed ValidAb™ |
| Cat No | HB11036 |
| Host | Goat |
| Clonality | Polyclonal |
| Target | Rabbit IgG H&L |
| Conjugate | Biotin |
| Description | Goat Anti-Rabbit IgG H&L (Biotin) preadsorbed secondary antibody. Part of the ValidAb™ range of highly validated, data-rich antibodies. |

Validation data



Product information

| | |
|----------------------|--|
| Immunogen | Purified rabbit IgG |
| Isotype | IgG |
| Purification | Immunogen affinity chromatography. Pre-adsorbed with mouse and human serum proteins |
| Concentration | 0.5 mg/ml |
| Formulation | Lyophilised. When reconstituted contains PBS with 1% recombinant albumin and less than 0.1% sodium azide |

Tested applications

| | |
|---|---|
| Applications | ELISA, ICC, IHC, IHC-P, WB, IHC(IF) |
| Western blot optimal concentration | Tested at 1:10,000. Optimise dependent upon assay. |
| IHC(IF) optimal concentration | Tested at 1:250 dilution (2µg/ml). Optimise dependent upon assay. |
| IHC-P optimal concentration | Tested at 1:300 dilution. Optimise dependent upon assay. |

ICC optimal concentration
Negative control

Tested at 1:250 dilution (2µg/ml). Optimise dependent upon assay.
While this antibody has been cross-adsorbed to reduce non-specific binding it is still often worthwhile to conduct a control experiment where the primary antibody is omitted to give confidence that the staining pattern observed is specific.

Storage & Handling

Storage instructions
Reconstitution advice

-20 °C then use reconstitution advice
We recommend reconstituting with either:

- dH₂O and storing at 4 °C
- 50:50 ratio of dH₂O to glycerol and storing at -20 °C
- dH₂O then aliquot and store at -80 °C

Take care when opening as the precipitate is extremely light and can easily be lost if disturbed. When reconstituting make sure that the antibody is thoroughly dissolved by pipetting up and down before giving the antibody a brief spin at <10,000g to make sure that all material is recovered and at the bottom of the tube.

Important

For more information please see our detailed guide on [storing and using your antibody](#)
This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

References

The use of antiavidin antibody and avidin-biotin-peroxidase complex in immunoperoxidase technics.

Hsu SM et al (1981) American journal of clinical pathology 75

PubMedID [6167159](#)

Antigen Retrieval and Signal Amplification.

Dunkenberger L et al (2022) Methods in molecular biology (Clifton, N.J.) 2422

PubMedID [34859399](#)
