


# DATASHEET

Streptavidin Janelia Fluor® 646

## Product overview

<b>Name</b>	Streptavidin Janelia Fluor® 646
<b>Cat No</b>	HB17045
<b>Biological description</b>	<p>Streptavidin Janelia Fluor® 646 is a biotin binding protein conjugated with the fluorescent dye Janelia Fluor® 646 and can be used to detect biotin labelled molecules such as nucleic acids, antibodies, and other proteins. Biotinylated antibodies are bound with extremely high affinity by Streptavidin Janelia Fluor® 646 enabling immunofluorescent detection in IHC, ICC, flow cytometry and Western blot. Janelia Fluor® 646 and the other members of the Janelia Fluor® family are bright and highly photostable fluorophores particularly suited for super resolution imaging such as dSTORM and STED.</p>
<b>Key features:</b>	<ul style="list-style-type: none"><li>• Conjugated with Janelia Fluor® 646 (Ex: 652nm, Em: 675nm)</li><li>• Supplied as a more stable lyophilate</li><li>• Bright and photostable signal for repeated imaging</li><li>• For use in IHC(IF), ICC, Western blotting and Flow cytometry</li><li>• Suited for super resolution imaging including dSTORM and STED</li></ul>
<b>Species of origin</b>	E. coli
<b>Applications</b>	fluorescence imaging, ICC, IF, IHC
<b>Description</b>	Janelia Fluor® 646 conjugated streptavidin for detection and signal amplification of biotin coupled proteins and antibodies.

## Images



## Biological Data

**Application notes**

**#Protocol 1: Detecting biotin-labelled antibodies in IHC**

1. Incubate free floating rat brain sections (40µm) in sodium borohydride (NaBH<sub>4</sub>) for 15 minutes followed by 2 hours in blocking buffer (0.05M glycine, 2% BSA and 3% donkey serum).
2. Incubate sections with primary antibody in blocking buffer at 4 °C overnight, as in our [IHC protocol](#).

3. Wash sections three times in PBST for 5 minutes each.
  4. Incubate sections with 2 µg/mL goat anti-mouse biotin antibody **HB11345** or goat anti-rabbit antibody **HB11036** diluted in blocking buffer for 2 hours at RT.
  5. Wash sections three times in PBST for 5 minutes each.
  6. Incubate sections with 1 µg/mL Streptavidin Janelia Fluor® 646 in blocking buffer for 2 hours.
  7. Wash sections three time in PBST for 5 minutes each.
  8. Incubate sections with 10 µg/mL DAPI for 10 minutes.
  9. Wash sections in dH<sub>2</sub>O, mount on glass slides with mounting media and cover with coverslip.
  10. Image the sections on a microscope using a 640nm laser or Cy5 filter set to excite Streptavidin Janelia Fluor® 646.
- 

## Solubility & Handling

**Storage instructions**  
**Reconstitution advice**

-20 °C then use reconstitution advice