

# DATASHEET

## Streptavidin Janelia Fluor® 525


### Product overview

<b>Name</b>	Streptavidin Janelia Fluor® 525
<b>Cat No</b>	HB15382
<b>Biological description</b>	Streptavidin Janelia Fluor® 525 is a biotin binding protein conjugated with the fluorescent dye Janelia Fluor® 525 and can be used to detect biotin labelled molecules such as nucleic acids, antibodies, and other proteins. Biotinylated antibodies are bound with extremely high affinity by Streptavidin Janelia Fluor® 525 enabling immunofluorescent detection in IHC, ICC, flow cytometry and Western blot. Janelia Fluor® 525 and the other members of the Janelia Fluor® family are bright and highly photostable fluorophores particularly suited for super resolution imaging such as dSTORM and STED.
<b>Species of origin</b>	E. coli
<b>Applications</b>	fluorescence imaging, ICC, IF, IHC
<b>Description</b>	Janelia Fluor® 525 conjugated streptavidin for detection and signal amplification of biotin coupled proteins and antibodies.

#### Key features:

- Conjugated with Janelia Fluor® 525 (Ex: 534nm, Em: 559nm)
- Supplied as a more stable lyophilate
- Bright and photostable signal for repeated imaging
- For use in IHC(IF), ICC, Western blotting and Flow cytometry
- Suited for super resolution imaging including dSTORM and STED

### Images



### Biological Data

<b>Application notes</b>	<b>#Protocol 1: Detecting biotin-labelled antibodies in IHC</b> <ol style="list-style-type: none"><li>1. Incubate free floating rat brain sections (40µm) in sodium borohydride (NaBH<sub>4</sub>) for 15 minutes followed by 2 hours in blocking buffer (0.05M glycine, 2% BSA and 3% donkey serum).</li><li>2. Incubate sections with primary antibody in blocking buffer at 4 °C overnight, as in our <a href="#">IHC protocol</a>.</li><li>3. Wash sections three times in PBST for 5 minutes each.</li><li>4. Incubate sections with 2 µg/mL goat anti-mouse biotin antibody <a href="#">HB11345</a> or goat anti-rabbit antibody <a href="#">HB11036</a> diluted in blocking buffer for 2 hours at RT.</li></ol>
--------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

5. Wash sections three times in PBST for 5 minutes each.
  6. Incubate sections with 1 µg/mL Streptavidin Janelia Fluor® 525 in blocking buffer for 2 hours.
  7. Wash sections three times in PBST for 5 minutes each.
  8. Incubate sections with 10 µg/mL DAPI for 10 minutes.
  9. Wash sections in dH<sub>2</sub>O, mount on glass slides with mounting media and cover with coverslip.
  10. Image the sections on a microscope using a 532nm laser or TRITC filter set to excite Streptavidin Janelia Fluor® 525.
- 

## Solubility & Handling

**Storage instructions**  
**Reconstitution advice**

-20 °C then use reconstitution advice