

# **Protocol Booklet**

Product Code(s) HB14861

Product Name(s) Fluo-4 AM Ultrapure (>98%)

**Purpose** Measurement of intracellular Ca<sup>2+</sup> in cultured cells

**Please note**: This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use



## **Contents**

Product Overview	3
Components & Storage	3
Protocol	3
Recipes	4
Fluo-4 AM Loading Solution	
HEPES-buffered Hank's Balanced Salt Solution (Assay Buffer)	4
Guidelines, precautions, troubleshooting	4
Contact	4
For customers in the UK, Europe and Rest of the World	4
For customers in the USA. Canada and South America	,



## **Product Overview**

Fluo-4 AM Ultrapure (>98%) is a frequently used single-wavelength  $Ca^{2+}$  indicator ( $Ca^{2+}$   $K_d$  = 0.35 $\mu$ M) that is membrane permeable and compatible with flow cytometry, plate reader, and fluorescence microscopy assays. Fluo-4 AM has excitation and emission peaks at 490nm and 515nm respectively which can be visualised using standard GFP microscope filters.

# **Components & Storage**

Fluo-4 AM is provided as either:

SKU	Component	Quantity	Storage Temperature
HB14861-500µg	Fluo-4 AM Ultrapure (>98%)	500µg	-20°C
HB14861-10x50μg	Fluo-4 AM Ultrapure (>98%)	10x50µg	-20°C

This protocol additionally requires:

Component	Quantity	Storage Temperature
DMSO	25µl	RT
Pluronic F-127	10mg	4°C
(optional) Probenecid	7.7mg	4°C
Assay Buffer (HEPES-buffered Hank's Balanced Salt Solution (pH = 7.3)*	10ml	RT

<sup>\*</sup> Please see recipe at the end of this protocol book.

## **Protocol**

The following protocol provides general guidelines for using Fluo-4 AM to measure intracellular calcium in cultured cells. All loading conditions (dye concentration, temperature, and time) should be optimized for your specific assay, application, and instrumentation.

- 1. Culture cells following standard protocols to approximately 80-100% confluence.
- 2. Prepare the loading solution freshly following the below table, vortex well and use within 2 hours.
- 3. Remove the cell culture medium, briefly wash in plain media (without serum), then add dye loading solution. Recommend volumes are:
  - a. 35mm dish / 6-well plate 1.5 mL/well,
  - b. 96 well plate 100 µL/well,
  - c. 384 well plate 20 µL/well,
- 4. Incubate in a cell culture incubator at 37°C for 60 minutes.
- 5. Read fluorescence using a plate reader (Excitation: 490nm, Emission 515nm) or image using a fluorescence microscope using filters for FITC/GFP



# **Recipes**

#### Fluo-4 AM Loading Solution

Component	Concentration	Quantity	Notes
Fluo-4 AM Ultrapure (>98%)	4.4µM	50µg	Dissolve in DMSO then aliquot and store any unused dye at -20°C
Assay Buffer	1X	10ml	Normally HEPES buffered HBSS but other buffers have been also successfully used.
Pluronic F-127	0.1%	10mg	Surfactant that helps the dissolution of dye therefore ensuring even dye distribution and cellular loading.
(optional) Probenecid	2.7mM	7.7mg	Anion transport inhibitor that improves intracellular dye retention. Not required for all cell types, it is recommended in most cases to optimize assay performance.

Please note: Combine components then vortex thoroughly. Use within 2 hours of creation. Do not freeze.

#### **HEPES-buffered Hank's Balanced Salt Solution (Assay Buffer)**

Component	MW (g/mol)	g/L	Concentration (mM)
Calcium Chloride	110.98	0.14	1.26
Magnesium Chloride Hexahydrate	203.30	0.1	0.49
Magnesium Sulfate Heptahydrate	246.47	0.1	0.41
Potassium Chloride	74.55	0.4	5.33
Potassium Phosphate Monobasic	136.09	0.06	0.44
Sodium Bicarbonate	84.01	0.35	4.17
Sodium Chloride	58.44	8	138.00
Sodium Phosphate Dibasic	141.96	0.048	0.34
D-Glucose (Dextrose)	180.16	1	5.56
HEPES	238.30	4.76	20.00

Please note: Add all components to dH<sub>2</sub>O, mix well then adjust to pH 7.3

# Guidelines, precautions, troubleshooting

Please contact our technical support team at technicalhelp@hellobio.com for advice on how to resolve any problems encountered when using this product. Observe safe laboratory practice and consult the safety datasheet. Please see the datasheet on our website for general guidelines, precautions, limitations on the use of the product.

#### **Contact**

For customers in the UK, Europe and Rest of the World

Customer

Customer Care <a href="mailto:customercare@hellobio.com">customercare@hellobio.com</a> customercare-usa@hellobio.com Care

For customers in the USA, Canada

and South America

Technical Technical technicalhelp@hellobio.com

technicalhelp@hellobio.com support support

By telephone: +44(0)117 318 0505 By telephone: +1-609-683-7500

+44(0)117 981 1601 +1-609-228-4994 By fax: By fax:

Opening Opening hours: 8.30 am - 5.00 pm GMT weekdays 9.00 am - 5.00 pm EST weekdays hours:

